



HOMEOPATHYRESEARCHINSTITUTE

Newsletter

Issue 3, January 2009

"To facilitate & disseminate high quality homeopathy research."

Welcome to the third newsletter of the Homeopathy Research Institute, a newsletter for everyone who wants to understand and support scientific research into homeopathy.

Homeopathy for women with menopausal symptoms: a review of the evidence

Introduction

Many women consult homeopaths for problems associated with the menopause e.g. hot flushes, night sweats, fatigue, anxiety, depression, mood swings, lack of confidence, vaginal dryness, and joint pain. A recent audit of patients receiving treatment from medically qualified homeopaths at five NHS homeopathic hospitals reported that the menopause was the third most common reason for patients to be referred for treatment (Thompson et al., 2008) and a study of professional homeopaths reported that 12.3 % of patients requested treatment for female complaints (Relton et al., 2006). This edition of the HRI newsletter summarises the existing evidence for the effectiveness of homeopathy in the treatment of menopausal symptoms. Two main types of evidence are reported, evidence from observational studies and case studies of treatment by homeopaths, and the more complex 'experimental' evidence from randomised controlled trials (RCTs) of 'homeopathy'.

Observational evidence of treatment by homeopaths

There is a significant amount of published observational evidence which reports the results of patients suffering from menopausal symptoms treated by homeopaths using individualised (classical) homeopathy. Individualised homeopathy consists of a series of in depth interviews with a strong focus on the patient's subjective experience to match the homeopathic medicine to the totality of symptoms (including any psychological symptoms) that emerge during a consultation and is regarded as the gold standard of homeopathic care.

Two observational studies report the outcomes of patients treated by homeopaths at NHS homeopathic hospitals (Clover & Ratsey, 2002; Thompson & Reilly, 2003) and an audit (Relton & Weatherley-Jones, 2005) reports the outcomes of patients in an NHS community menopause clinic.

All patients sought homeopathic treatment for one or more of the following menopausal symptoms: hot flushes, vaginal dryness, mood disturbance, fatigue. Many women receiving treatment for breast cancer suffer from menopausal symptoms such as hot flushes. One study reports the results of 45 patients who have had a diagnosis of breast cancer 35 of whom were suffering from hot flushes (Thompson & Reilly, 2003). Clover & Ratsey, describe the outcomes of 31 patients with hot flushes, the majority of whom have had a diagnosis of past or current breast cancer (20/31) and who also reported taking a wide range of medication (tamoxifen, HRT, antidepressants, clonidine, and chemotherapy). These two studies used patient self-assessment as their primary outcome. Clinically significant improvements were reported by Clover & Ratsey for hot flush frequency and severity, while Thompson & Reilly reported clinically significant improvements in effect of symptoms on daily living, mood, and quality of life. Finally, an audit of 102 women attending an NHS community menopause clinic demonstrated significant improvements in headaches, vasomotor symptoms, emotional and psychological symptoms, tiredness and fatigue (Relton & Weatherley-Jones, 2004).

Randomised controlled trials of homeopathic remedies

There have been four RCTs of homeopathic remedies conducted. Two small (i.e. 5 patients in total) randomised studies evaluated the use of homeopathy for menopausal symptoms in the climacteric (Bekkering, 1993; Gautier, 1983) but neither study was large enough to make the detection of a statistically significant difference between the groups possible.

Two full sized double blind placebo-controlled trials were conducted in hospital settings, one in the UK (Thompson et al, 2005) and the other in the USA (Jacobs et al., 2005).

The duration of the intervention varied between 16 weeks (Thompson et al., 2005) and 6 – 12 months (Jacobs et al., 2005). The patient sample sizes were 83 (Jacobs et al., 2005) and 53 (Thompson et al., 2005). Both RCTs used repeated consultations with a homeopath with either an individualised homeopathic remedy or placebo, however, Jacobs et al., (2005) had an additional treatment arm of a formulaic complex homeopathic remedy. Inclusion criteria for both trials were: three or more hot flushes a day and a history of breast cancer.

The patient mean age was 52 (Thompson et al., 2005) and 55 (Jacobs et al., 2005) and use of Tamoxifen was high (80% Thompson et al., 60% Jacobs et al.). Jacobs et al. reported a high dropout rate (28/83) perhaps due to the greater length and older age group. There was evidence that women taking the formulaic complex homeopathic remedy, containing homeopathic preparations of Sanguinaria, Amyl nitrate and Lachesis, experienced side effects with the development of new or proving symptoms including a significant increase in headaches.

Neither RCT found a statistically significant improvement in the primary outcome measures for 'homeopathy' over placebo. The study by Jacobs in 2005 did however show a positive trend for homeopathy in the reduction of hot flashes during the first three months ($p=0.1$) and a reduction in the Kupperman Menopausal Index ($p=0.1$) at one year. It is impossible to conclude whether these negative results are due to insufficient sample sizes, faulty trial designs or the ineffectiveness of the interventions.

Conclusions

None of the randomised controlled trials (RCTs) performed to date have reported a statistically significant effect for the use of homeopathic remedies in the treatment of menopausal symptoms. On the other hand, the observational evidence for the effectiveness of the treatment by a homeopath is overwhelmingly positive; suggesting it is a clinically relevant option in this clinical setting. In this instance, we must remember that the lack of evidence of effect reported by RCTs does not constitute evidence that there is no effect. Further well conducted trials of sufficient size are required to evaluate the clinical and cost effectiveness of treatment by a homeopath.

References

- Bekkering GM van den Bosch. (1993) Bedriegt Schone Schijn? Een Onderzoek Om De Gerapporteerde Werking Van Een Homeopathisch Middel Te Objectiveren. Huisarts en Wetenschap 36, 414-415.
- Clover, A., Ratsey, D., (2002), 'Homeopathic treatment of hot flushes: a pilot study', Homeopathy 91, 75-79.
- Gautier J. (1983), Essai Therapeutique Comparatif De L'action De La Clonidine et Du Lachesis Mutans Dans Le Traitement des Bouffes et De La Chaleur De La Menopause. Universite de Bordeaux (thesis).
- Jacobs, J., Herman P., Herron, K. et al. (2005), 'Homeopathy for menopausal symptoms in breast cancer survivors: a preliminary randomized controlled trial', Journal of Alternative & Complementary Medicine, 11, 1, 21-27.
- Relton, C., Chatfield, K., Partington, H., et al. (2007), 'Patients treated by homeopaths registered with the Society of Homeopaths: a pilot study'. Homeopathy, 96, 87-89.
- Relton, C., Weatherley-Jones, E. (2005), 'Homeopathy service in a NHS community menopause clinic: audit of clinical outcomes', Journal of the British Menopause society, 11, 2, 72-3.
- Thompson, E.A., Reilly, D., (2002), 'The homeopathic approach to symptom control in the cancer patient: a prospective observational study'. Palliative Medicine 16, 3, 227-33.
- Thompson, E.A., Reilly, D. (2003), 'The homeopathic approach to the treatment of symptoms of oestrogen withdrawal in breast cancer patients: A prospective observational study' Homeopathy, 92,131-134.
- Thompson, E.A., Montgomery, A., Douglas, D., Reilly, D. (2005), 'A pilot randomised double blind placebo-controlled trial of individualised homeopathy for symptoms of oestrogen withdrawal in breast cancer survivors'. Journal of Alternative & Complementary Medicine, 11,1,: 13-20
- Thompson, E.A., Mathie, R.T., Baitson, E.S., et al. (2008), 'Towards standard setting for patient-reported outcomes in the NHS homeopathic hospitals' Homeopathy, 97,114-121.

Author: C. Relton

The HRI needs your support

HRI is currently seeking funding and support from those with an interest in homeopathy research. If you want to subscribe to this newsletter or make a donation to the HRI, please go to our website.

Visit us online at www.homeoinst.org