

Examples of trials with less than 150 participants funded by NHMRC

Trial title	Trial URL	Brief description	Trial end year	Sample size
Does Aerobic or Resistance Exercise Training Improve Walking Ability in Chronic Stroke Patients?	clinicaltrials.gov/show/NCT00107068	Randomized, double blind; bio-equivalence study	2004	48
Phase II/III Multicentre Randomised Clinical Trial of Laser Treatment Plus 4 mg Intravitreal Triamcinolone Injection to Reduce Diabetic Macular Oedema	clinicaltrials.gov/show/NCT00148265	Randomized, double blind; phase 2, phase 3	2009	54
A Multicentre Randomised Clinical Trial of Intravitreal Bevacizumab (Avastin®) Versus Intravitreal Dexamethasone (Ozurdex™) for Persistent Diabetic Macular Oedema	clinicaltrials.gov/show/NCT01298076	Comparing two treatments, safety, randomized, single blind; Phase 2	2014	61
An Open Label Extension of the Phase II/III Clinical Trial of Intravitreal Triamcinolone on the Effects and Safety of Clinically Significant Diabetic Macular Oedema That Persists After Laser Treatment	clinicaltrials.gov/show/NCT00148330	Non-randomized, open label; phase 2, phase 3	2008	64
A Sham-controlled Study of Transcranial Direct Current Stimulation (tDCS) as a Treatment for Depression	clinicaltrials.gov/show/NCT00763230	Randomized, double blind; Phase 2, phase 3	2011	64
A Randomised Trial of Radical Chemo/Radiotherapy vs Radiotherapy Alone in the Definitive Management of Localised Muscle Invasive TCC of the Urinary Bladder	clinicaltrials.gov/show/NCT00330499	Randomized, open label, phase 3, efficacy	2010	67
Renal Denervation for Resistant Hypertension	clinicaltrials.gov/show/NCT01865240	Randomized, open label; Phase 4	2015	100
A RCT of Power Training and Treadmill Training to Improve Walking Ability in Sub-acute Stroke Patients	clinicaltrials.gov/show/NCT00108030	Randomized, single blind; phase 2, phase 3	2006	102